

Healthcare Personnel Attire and Devices as Fomites: A Systematic Review

Nicholas Haun, MD;¹ Christopher Hooper-Lane, MA;² Nasia Safdar, MD, PhD^{3,4}

BACKGROUND. Transmission of pathogens within the hospital environment remains a hazard for hospitalized patients. Healthcare personnel clothing and devices carried by them may harbor pathogens and contribute to the risk of pathogen transmission.

OBJECTIVE. To examine bacterial contamination of healthcare personnel attire and commonly used devices.

METHODS. Systematic review.

RESULTS. Of 1,175 studies screened, 72 individual studies assessed contamination of a variety of items, including white coats, neckties, stethoscopes, and mobile electronic devices, with varied pathogens including *Staphylococcus aureus*, including methicillin-resistant *S. aureus*, gram-negative rods, and enterococci. Contamination rates varied significantly across studies and by device but in general ranged from 0 to 32% for methicillin-resistant *S. aureus* and gram-negative rods. *Enterococcus* was a less common contaminant. Few studies explicitly evaluated for the presence of *Clostridium difficile*. Sampling and microbiologic techniques varied significantly across studies. Four studies evaluated for possible connection between healthcare personnel contaminants and clinical isolates with no unequivocally direct link identified.

CONCLUSIONS. Further studies to explore the relationship between healthcare personnel attire and devices and clinical infection are needed.

Infect Control Hosp Epidemiol 2016;1–7

Transmission of pathogens within the hospital environment remains a hazard for hospitalized patients. Organisms such as *Staphylococcus aureus*, including methicillin-resistant *S. aureus* (MRSA), and vancomycin-resistant enterococci are associated with considerable morbidity, mortality, and healthcare costs and can be transmitted via environmental surfaces and inanimate objects. Healthcare personnel (HCP) themselves may represent a mobile surface for transmission via their contaminated apparel. Efforts to improve hand hygiene and reforms such as the United Kingdom's dress code policy of "bare below the elbow" have attempted to reduce this risk, but the professional wardrobe and numerous devices carried by care providers still represent potential risks. However, the magnitude of the risk is unclear. We performed a systematic review of the literature to evaluate the bacterial contamination of HCP attire and commonly used devices.

METHODS

We undertook a systematic search for studies that assessed the prevalence of pathogenic bacterial contamination of apparel and devices carried by HCP. MEDLINE, Cumulative Index to

Nursing and Allied Health Literature, and Cochrane databases were searched. The following search terms were developed in MEDLINE and adapted for use in other databases: ("fomites"[MeSH] OR fomite* OR "Cross infection"[MeSH] OR nosocomial OR "Bacteria"[MeSH] OR "Bacterial Infections"[MeSH]) AND ("Equipment Contamination"[MeSH] OR "mobile phone" OR "mobile phones" OR "Cell Phones"[MeSH] OR "cellular phones" OR "cellular phone" OR Pager OR pagers OR Pens OR "writing utensil" OR "Personal Digital Assistant" OR "personal digital assistants" OR "Computers, Handheld"[MeSH] OR "smart device" OR "smart devices" OR ipad OR ipads OR purse OR purses OR handbag* OR badge OR badges OR lanyard* OR necktie* OR "white coat" OR "white coats" OR clothing OR uniforms OR attire OR stethoscope* OR otoscope* OR sphygmomanometer*) AND (health personnel OR physician OR physicians OR nurse OR nurses OR doctor OR doctors OR student OR students OR medical personnel). Related citations and bibliographies were also reviewed for additional studies of relevance. The search was last performed February 10, 2015. Studies were included if the prevalence of pathogenic bacteria, particularly *S. aureus* or gram-negative rods (GNR), was explicitly stated or able to be

Affiliations: 1. Division of Hospital Medicine, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin; 2. Ebling Library, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin; 3. Division of Infectious Disease, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin; 4. William S. Middleton Memorial Veterans Hospital, Madison, Wisconsin.

Received March 9, 2016; accepted July 7, 2016

© 2016 by The Society for Healthcare Epidemiology of America. All rights reserved. DOI: 10.1017/ice.2016.192

extracted, the study was published in 1995 or later, and it was available in English. Studies of contamination of hands, gloves, isolation gowns, or environment were excluded. Studies of fungal or viral contamination were also excluded.

Extracted data included study location, population, item studied, and prevalence of contamination. Gram-negative rod contamination was reported on the basis of the individual authors' description of which isolates are pathogenic. Results from studies that included both inpatient and outpatient HCP were pooled into a single combined prevalence. In studies that compared personal equipment with environmental (dedicated) equipment, only the prevalence of contamination of personal equipment was included. Only contamination data from pretreatment or controls were used from studies that tested sanitation strategies. Prevalence of hand contamination was not included.

RESULTS

The systematic search yielded a total of 1,175 studies, 115 of which met criteria for full review. Of these, along with additional review of relevant citations and references, 72 unique studies were identified as meeting search criteria. These studies are described in Table 1.^{1–76} Eighteen studies originated in the United States whereas the remainder were from Asia (24), Europe (19), Africa (5), other North American countries (5), and Australia (1). Various sampling techniques, microbiologic processes, and sensitivity tests were performed in the studies. Sampling techniques differed with the item studied, with 94% of phone studies using a sampling method using swabs whereas 60% of clothing studies used direct inoculation onto solid culture media. The most frequent microbiologic method was nonselective solid culture medium, such as blood agar, with or without additional selective media based on that study's pathogen of interest. Twenty-four studies analyzed contamination of stethoscopes, with MRSA contamination prevalence of 0–42% and GNR prevalence of 0–31%. Twenty-eight studies analyzed digital communication devices; 21 of these evaluated mobile phones explicitly. The range of MRSA contamination for phones was 0–20% and the range of GNR contamination for phones was 0–75%. One study¹ of tablets had MRSA contamination of 50%. Eight studies on white coats yielded rates of MRSA contamination of 0–16%, with one outlier² that was performed in the midst of an outbreak. GNR contamination of white coats ranged from 0 to 42%. Neckties had a reported MRSA contamination rate of 3%–32% and GNR contamination of 11%–23% in 5 studies. There was considerable variation in which areas of the white coats were sampled across the included studies.

Few studies explicitly evaluated for the presence of *Clostridium difficile*. One study³ directed at *C. difficile* contamination of stethoscopes identified a contamination rate of 5%, whereas a second⁴ identified no such contamination. Studies used a wide variety of classification schemes to report gram-negative rods, depending on the microbiologic methods used

for isolation. Some provided species-level data whereas others reported only “GNR,” “nonfermenting GNR,” or “coliforms.” *Enterococcus* contamination was inconsistently reported, and where it was included, vancomycin resistance was rare. The exception is a study⁵ of nurses' uniforms with 39% contamination with vancomycin-resistant enterococci. Because of high variation in sampling technique and equipment, microbiologic methods, and reporting, no attempt was made to pool data from all studies or conduct a meta-analysis.

Three studies^{5–7} were prospective. The remaining studies were cross-sectional. Four studies^{2,8–10} attempted to correlate device or apparel contamination with clinical isolates.

DISCUSSION

We found that stethoscopes, digital devices, white coats, and neckties are commonly contaminated with bacterial pathogens including *S. aureus* (including MRSA) and GNRs, though there was high interfacility and interstudy variability. This may be due in part to the varied clinical settings included—inpatient vs outpatient vs emergency room and adult vs pediatric patient populations. However, even within a particular setting, variability persists. Possibilities include variable endemic rates of MRSA in the patient population and the hospital environment or local differences in hand hygiene or cleaning practices that may confound attire contamination. Another possibility is that there is no standardized approach regarding how to sample attire and devices, there may be differences in the ability of different types of swabs to pick up pathogens, and there may be variable efficiency of transfer of pathogens to and from different materials. For all these studies, sampling was not performed longitudinally, thus limiting the ability to evaluate persistent presence of pathogens. For most studies, the cleaning of devices or attire was not reported or taken into account at the time of sampling, which also may explain the wide ranges of prevalence of pathogens recovered from the items under study.

This review expands upon the findings of a prior systematic review¹⁰ that focused only on contamination of digital devices and included 15 studies that sampled mobile phones, pagers, and personal data assistants for contamination with pathogens including MRSA and GNRs.

There is no evidence to date directly linking HCP-borne fomites with patient infection other than a report¹¹ of sternal wound infections linked to a nurse anesthetist. In that case, a cluster of 3 sternal wound infections due to *Gordonia bronchialis* triggered an epidemiologic survey that identified a nurse anesthetist involved in all 3 cases as the likely link. *Gordonia* was isolated from the nurse's axilla and hands as well as her scrubs and purse. Pulsed-field gel electrophoresis confirmed relatedness of these strains with clinical samples in each case. The pathogen was also identified in her home, and the authors suspect that home-laundering of scrubs may have led to the contamination. Our study does not attempt to correlate the presence of organisms on the objects sampled with

TABLE 1. Systematic Review of Prior Studies on Healthcare Personnel Attire and Devices as Fomites, Sorted by Item, With Reported Prevalence of Contamination With Various Pathogens

Source	Population	Sample	<i>S. aureus</i>	MRSA	GNR	Notes
Stethoscopes						
Alleyne et al ³	Inpatient physicians, UK	Stethoscopes	NR	NR	NR	5% (3/61) <i>C. difficile</i>
Bandi et al ¹³	Inpatient pediatric HCP, UK	Stethoscopes	3% (1/40)	0% (0/40)	3% (1/40)	
Bernard et al ¹⁴	Inpatient HCP, France	Stethoscopes	4% (15/355)	0% (1/355)	7% (25/355)	No MDR GNR
Burkharie et al ¹⁵	Inpatient HCP, Saudi Arabia	Stethoscopes	0% (0/100)	0% (0/100)	9% (9/100)	
Campos-Murquia et al ¹⁶	Inpatient HCP, Mexico	Stethoscopes	38% (43/112)	16% (18/112)	3% (3/112)	4% (4/112) <i>Enterococcus</i>
Cohen et al ¹⁷	Outpatient pediatric physicians, Israel	Stethoscopes	55% (30/55)	7% (4/55)	18% (10/55)	
Faffiori et al ¹⁸	ED physicians, Greece	Stethoscopes	7% (6/88)	2% (2/88)	3% (3/88)	
Fenelon et al ¹⁹	Inpatient HCP, Ireland	Stethoscopes	NR	0% (0/44)	NR	
Jones et al ²⁰	ED HCP, USA	Stethoscopes	17% (25/150)	NR	NR	
Marinella et al ⁴	Inpatient HCP, USA	Stethoscopes	38% (15/40)	NR	5% (2/40)	0% (0/40) <i>C. difficile</i> 5% (2/40) <i>Enterococcus</i>
Merlin et al ²¹	EMS HCP, USA	Stethoscopes	NR	32% (16/50)	NR	
Nunez et al ²²	ED HCP, Spain	Stethoscopes	6% (7/122)	0% (0/122)	5% (6/122)	
Pandey et al ²³	Hospital physicians, USA	Stethoscopes	28% (22/80)	NR	18% (14/80)	
Panhotra et al ²⁴	Inpatient physicians, Saudi Arabia	Stethoscopes	48% (23/48)	4% (2/48)	15% (7/48)	8% (4/48) MDR <i>Pseudomonas</i>
Russell et al ²⁵	Inpatient HCP, USA	Stethoscopes	NR	0% (0/141)	NR	
Schroeder et al ²⁶	Inpatient and outpatient HCP, USA	Stethoscopes	5% (5/92)	3% (3/92)	18% (17/92)	
Sengupta, et al ²⁷	Inpatient pediatrics, India	Stethoscopes	28% (12/43)	28% (12/43)	21% (9/43) 5% (2/43) FQR	5% (2/42) <i>Enterococcus</i>
Smith et al ²⁸	Inpatient and outpatient HCP, USA	Stethoscopes	12% (24/200)	2% (4/200)	1% (2/200)	
Shiferaw et al ²⁹	Inpatient HCP, Ethiopia	Stethoscopes	45% (79/176)	12% (21/176)	31% (54/176)	
Sood et al ³⁰	Inpatient physicians, India	Stethoscopes	18% (19/106)	4% (4/106)	0% (0/106)	5% (5/106) <i>Enterococcus</i>
Tang et al ³¹	ED staff, Canada	Stethoscopes	1% (1/100)	0% (0/100)	12% (12/100)	
Uneke et al ³²	HCP, Nigeria	Stethoscopes	42% (45/107)	42% (45/107)	25% (27/107), 17% (18/107) MDR	11% (12/107) <i>Enterococcus</i>
Whittington et al ³³	ICU HCP, UK	Stethoscopes	9% (2/22)	5% (1/22)	14% (3/22)	
Youngster et al ³⁴	Inpatient pediatric physicians, Israel	Stethoscopes	9% (4/43)	2% (1/43)	21% (9/43)	
Digital devices						
Akinyemi et al ³⁵	Hospital HCP, Nigeria	Mobile phones	16% (14/90)	0% (0/90)	9% (8/90)	4% (4/90) <i>Enterococcus</i>
Beer et al ³⁶	Inpatient HCP, Canada	Pagers	10% (10/100)	1% (1/100)	4% (4/100)	
Borer et al ³⁷	Inpatient HCP, Israel	Mobile phones	NR	NR	12% (15/124) <i>Acinetobacter</i>	
Braddy et al ³⁸	Inpatient HCP, USA	PDA's	2% (2/82)	0% (0/82)	2% (2/82)	1% (1/78) anaerobe
Brady et al ³⁹	Operating rooms, UK	Phones, pagers, PDA's	4% (3/78)	0% (0/78)	6% (5/78)	
Brady et al ⁴⁰	Inpatient HCP, UK	Mobile phones	8% (8/105)	2% (2/105)	5% (5/105)	1% (1/105) <i>Enterococcus</i>
Datta et al ⁴¹	HCP, India	Mobile phones	36% (72/200)	13% (26/200)	0% (0/200)	0% (0/200) <i>Enterococcus</i>
Goldblatt et al ⁴²	Inpatient HCP, Israel and USA	Mobile phones	4% (17/400)	3% (10/400)	17% (67/400)	
Hassoun et al ⁴³	Inpatient HCP, USA	PDA's	11% (8/75)	8% (6/75)	0% (0/75)	1% (1/75) VRE
Hirsch et al ¹	Inpatient and outpatient pharmacy faculty, USA	Tablets	23% (7/30)	50% (15/30)	7% (2/30)	3% (1/30) VRE
Jayalakshmi et al ⁴⁴	Clinical and non-clinical physicians, India	Mobile phones	23% (33/144)	3% (4/144)	6% (9/144)	1% (1/144) <i>Enterococcus</i>
Karabay et al ⁴⁵	Inpatient HCP, Turkey	Mobile phones	7% (9/122)	0% (0/122)	7% (8/122)	2% (2/122) VSE 0% VRE
Khivsara et al ⁹	HCP, India	Mobile phones	40% (12/30)	10% (3/30)	NR	
Kilic et al ⁴⁶	Inpatient HCP, Pakistan	Mobile phones	8% (8/106)	NR	1% (1/106)	
Lee et al ⁴⁷	Inpatient HCP, South Korea	Mobile phones	25% (50/203)	4% (8/203)	4% (8/203)	0% (1/203) <i>Enterococcus</i>
Namias et al ⁴⁸	Inpatient HCP, USA	Pagers	19% (7/36)	NR	22% (8/36)	6% (2/36) <i>Enterococcus</i>
Nwankwo et al ⁴⁹	Hospital HCP, Nigeria	Mobile phones	25% (14/56)	NR	59% (33/56)	High prevalence of antibiotic-resistance in GNR
Pandey et al ²³	Hospital physicians, USA	Mobile phones	8% (10/126)	NR	19% (24/126)	
Ramesh et al ⁵⁰	Inpatient HCP, Barbados	Mobile phones	0% (0/101)	0% (0/101)	15% (15/101)	
Sadat-Ali et al ⁵¹	Inpatient and outpatient HCP, Saudi Arabia	Mobile phones	15% (44/288)	3% (8/288)	9% (26/288)	3% (10/288) <i>Enterococcus</i>
Saxena et al ⁵²	Physicians and nurses, India	Mobile phones	26% (26/100)	17% (17/100)	2% (2/100)	
Singh et al ⁵³	Inpatient and outpatient HCP, USA	Pagers	21% (21/100)	3% (3/100)	0% (0/100)	0% (0/100) <i>Enterococcus</i>
Smith et al ⁵⁴	Inpatient physicians, USA	Notebook computers	2% (1/60)	0% (0/60)	2% (1/60)	No <i>C. diff</i> recovered on culture of 17 devices; not tested for spores
Srikanth et al ⁵⁵	HCP, India	Mobile phones	12% (6/51)	4% (2/51)	18% (9/51)	
Tambekar et al ⁵⁶	Physicians, India	Mobile phones	24% (18/75)	20% (15/75)	75% (56/75)	

TABLE 1. *Continued*

Source	Population	Sample	<i>S. aureus</i>	MRSA	GNR	Notes
Ulger et al ⁵⁷	OR and ICU HCP, Turkey	Mobile phones	25% (50/200)	13% (26/200)	8% (15/200) coliforms 10% (19/200) NFGN	4% (7/200) <i>Enterococcus</i>
Ustun et al ⁵⁸	Inpatient HCP, Turkey	Mobile phones	25% (45/183)	9% (17/183)	23% (42/183)	11% ESBL + GNR 1% (1/183) <i>Enterococcus</i>
Walia et al ⁵⁹	Inpatient, outpatient, and dental HCP, India	Mobile phones	18% (54/300)	11% (33/300)	13% (39/300)	
White coats						
Burden et al ⁶	Inpatient physicians, USA	White coats	NR	16% (8/50)	NR	
Loh et al ⁶⁰	Medical students, UK	White coats	5% (5/100)	0% (0/100)	GNR isolated on 3%, not deemed pathogenic	
Munoz-Price et al ⁶¹	ICU HCP, USA	White coats	32% (7/22)	0% (0/22)	32% (7/22)	5% (1/22) <i>Enterococcus</i>
Osawa et al ²	Inpatient HCP, Japan	White coats	NR	79% (11/14); later 38% (9/24)		Performed during and after MRSA outbreak
Pandey et al ²³	Hospital physicians, USA	White coat	6% (8/130)	NR	NR	
Treacle et al ⁶²	Grand rounds attendees, USA	White coats	23% (34/149)	4% (6/149)	NR	0% VRE
Uneke et al ⁶³	Physicians, Nigeria	White coats	17% (18/103)	NR	26% (27/103)	
Wiener-Well et al ⁶⁴	Inpatient HCP, Israel	White coats	19% (10/52)	NR	42% (22/52)	<i>Acinetobacter</i> predominates GNRs. Rates reported per no. of cultures positive. Did not report contamination per coat.
Neckties						
Ditchburn ⁶⁵	Hospital physicians, UK	Neckties	20% (8/40)	3% (1/40)	NR	
Koh et al ⁶⁶	Inpatient physicians, Malaysia	Neckties	52% (26/50)	32% (16/50)	NR	
Lopez et al ⁶⁷	Physicians, UK	Neckties	32% (16/50)	NR	NR	
McGovern et al ⁶⁸	Physicians, Scotland	Neckties	11% (10/95)	8% (8/95)	11% (10/95)	0% VRE
Steinlechner et al ⁸	Orthopedic surgeons, UK	Neckties	8% (2/26)	NR	23% (6/26)	
Pens						
Bhat et al ⁶⁹	ICU physicians and nurses, India	Pens	8% (6/75)	3% (2/75)	0% (0/75)	
Datz et al ⁷⁰	Inpatient physicians, Austria	Pens	2% (1/42)	0% (0/42)	7% (3/42)	
French et al ⁷	Inpatient HCP, UK	Pens	NR	25% (9/36)	0% (0/8) MDR <i>Klebsiella</i>	17% (1/6) VRE Items sampled from wards with outbreaks with that pathogen.
Halton et al ⁷¹	Inpatient HCP, USA	Pens	NR	NR	31% (4/13)	38% (5/13) <i>Staphylococcus</i> , not speciated
Pandey et al ²³	Hospital physicians, USA	Pens	14% (14/100)	NR	NR	
Wolfe et al ⁷²	ICU RTs, USA	Pens	0% (0/20)	0% (0/20)	0% (0/20)	
Other apparel						
Burden et al ⁶	Inpatient physicians, USA	Short-sleeved uniforms	NR	20% (10/50)	NR	
Feldman et al ⁷³	Female physicians, USA	Purses	0% (0/13)	0% (0/13)	NR	69% (9/13) skin flora
Gaspard et al ⁷⁴	Long-term care facility HCP, France	Uniforms	NR	17% (43/256) waist 16% (42/ 256) pocket	NR	
Kotsanas et al ⁷⁵	Inpatient HCP, Australia	ID badge and lanyard	19% (11/59) badges, 25% (15/59) lanyards	5% (3/59) badges, 7% (4/59) lanyards	0% (0/59) badges, 14% (8/59) lanyards	2% (1/59) lanyards with <i>Enterococcus</i>
Munoz-Price et al ⁶¹	ICU HCP, USA	Scrubs	11% (11/97)	4% (4/97)	11% (11/97)	3% (3/97) <i>Enterococcus</i>
Ota et al ⁷⁶	Inpatient HCP, Canada	ID badge	7% (8/118)	0% (0/118)	6% (7/118)	
Perry et al ⁵	Nurses, UK	Uniforms	NR	14% (8/57)	NR	39% (22/57) VRE; 19% (11/57) <i>C. difficile</i>
Saxena et al ⁵²	Physicians and nurses, India	Rings	50% (50/100)	28% (28/100)	32% (32/100)	
Wiener-Well et al ⁶⁴	Inpatient HCP, Israel	White coats	22% (32/147)	5% (8/147)	78% (115/147)	60% <i>Acinetobacter</i>

NOTE. Prevalence is reported as % of total items contaminated followed by number over *n* in parentheses; "*S. aureus*" indicates total contamination by methicillin-susceptible *Staphylococcus aureus* and methicillin-resistant *S. aureus* (MRSA), and "GNR" indicates pathogenic gram-negative rods as differentiated by the author. *C. difficile*, *Clostridium difficile*; ED, emergency department; EMS, emergency medical services; ESBL, extended-spectrum beta-lactamase; FQR, fluoroquinolone-resistant; HCP, healthcare personnel; ICU, intensive care unit; MDR, multidrug-resistant; NFGN, nonfermenting gram-negative rods; NR, not reported; PDA, personal data assistant; OR, operating room; RT, respiratory therapist; VRE, vancomycin-resistant *Enterococcus*; VSE, vancomycin-sensitive *Enterococcus*.

transmission to patients or clinical infection, though some individual studies did attempt to make this link. Steinlechner et al⁸ found 45% correspondence at the species level only between clinical isolates from surgical wound infections on inpatient orthopedic wards and isolates contaminating neckties from orthopedic surgeons. French et al⁷ found that MRSA isolates from HCP pens had antimicrobial resistance patterns that corresponded to clinical isolates from an ongoing outbreak. Osawa et al² found that isolates of MRSA contaminating white coats during an outbreak on an inpatient ward were not genetically similar (on the basis of pulsed-field gel electrophoresis) to the outbreak strain whereas a later sampling in a non-outbreak setting did indicate genetic relatedness of clinical and HCP-derived strains. Khivsara et al⁹ found that although antibiotic sensitivities were identical among MRSA isolated from HCP mobile phones and clinically derived specimens, molecular typing indicated that the strains were not related.

In 2014, the Society for Healthcare Epidemiology of America published recommendations¹² for healthcare facilities to address HCP attire, including the consideration of “bare below the elbows” policies, provision of white coat laundering services, and provision of hooks to remove white coats prior to patient contact. However, there remains a paucity of data linking attire or device contamination with patient infection, and the findings of this review call for research in the area of attire and mobile and other devices used by HCP.

Our findings have implications for clinicians and infection preventionists. Once hand hygiene practices have been optimized, attention to reducing reservoirs of organisms that may exist in clothing and devices is a reasonable next step in infection control. Possibilities include incorporation of attire policies consistent with Society for Healthcare Epidemiology of America recommendations, inclusion of stethoscope cleaning as part of hand hygiene practices, and implementation and enforcement of policies for cleaning shared patient items on a schedule agreed upon by unit staff.

Our study has limitations. The first limitation was the variability of methods in the individual studies. The included studies varied significantly by methods of sampling, including both site and method (eg, use of swab versus direct inoculation onto culture media). The microbiologic evaluation used also varied in many facets, including the extent to which pathogens were isolated and speciated and tested for antibiotic sensitivity.

Next steps for research in this area include the development of standardized methods and protocols that would enable more meaningful comparison between studies and institutions. A serial sampling strategy using longitudinal study design may yield important insights into the persistence of bacterial contamination. Given the paucity of data regarding *C. difficile* contamination relative to the importance of this pathogen in healthcare-associated infection in this era, further study specific to this pathogen is essential. Finally, while the use of new technology such as antimicrobial-impregnated fabrics or accessories has recently gained ground,

methodologically rigorous study designs are needed to evaluate the impact of this novel technology on clinical outcomes rather than solely focusing on reducing contamination.

ACKNOWLEDGMENTS

Financial support. Department of Veterans Affairs, Veterans Health Administration, Health Services Research and Development Service Quality Enhancement Research Initiative (project no. PEC 15-248); and the Veterans Health Administration National Center for Patient Safety Center of Inquiry, United States Department of Veterans Affairs.

Potential conflicts of interest. All authors report no conflicts of interest relevant to this article.

Disclaimer: The views expressed in this article are those of the author(s) and do not necessarily represent the views of the Department of Veterans Affairs.

Address correspondence to Nasia Safdar, MD, PhD, 5138 MFCB, 1685 Highland Ave, Madison, WI 53705 (ns2@medicine.wisc.edu).

REFERENCES

- Hirsch EB, Raux BR, Lancaster JW, Mann RL, Leonard SN. Surface microbiology of the iPad tablet computer and the potential to serve as a fomite in both inpatient practice settings as well as outside of the hospital environment. *PLOS ONE* 2014;9: e111250.
- Osawa K, Baba C, Ishimoto T, et al. Significance of methicillin-resistant *Staphylococcus aureus* (MRSA) survey in a university teaching hospital. *J Infect Chemother* 2003;9:172–177.
- Alleyne SA, Hussain AM, Clokie M, Jenkins DR. Stethoscopes: potential vectors of *Clostridium difficile*. *J Hosp Infect* 2009; 73:187–189.
- Marinella MA, Pierson C, Chenoweth C. The stethoscope: a potential source of nosocomial infection? *Arch Intern Med* 1997;157:786–790.
- Perry C, Marshall R, Jones E. Bacterial contamination of uniforms. *J Hosp Infect* 2001;48:238–241.
- Burden M, Cervantes L, Weed D, Keniston A, Price CS, Albert RK. Newly cleaned physician uniforms and infrequently washed white coats have similar rates of bacterial contamination after an 8-hour workday: a randomized controlled trial. *J Hosp Med* 2011;6:177–182.
- French G, Rayner D, Branson M, Walsh M. Contamination of doctors' and nurses' pens with nosocomial pathogens. *Lancet Lond Engl* 1998;351:213.
- Steinlechner C, Wilding G, Cumberland N. Microbes on ties: do they correlate with wound infection? *Bull R Coll Surg Engl* 2002;84:307–309.
- Khivsara A, Sushma T, Dahashree B. Typing of *Staphylococcus aureus* from mobile phones and clinical samples. *Curr Sci* 2006; 90:910–912.
- Brady RRW, Verran J, Damani NN, Gibb AP. Review of mobile communication devices as potential reservoirs of nosocomial pathogens. *J Hosp Infect* 2009;71:295–300.
- Wright SN, Gerry JS, Busowski MT, et al. *Gordonia bronchialis* sternal wound infection in 3 patients following open heart surgery: intraoperative transmission from a healthcare worker. *Infect Control Hosp Epidemiol* 2012;33:1238–1241.
- Bearman G, Bryant K, Leekha S, et al. Healthcare Personnel attire in non-operating-room settings. *Infect Control Hosp Epidemiol* 2014;35:107–121.

13. Bandi S, Uddin L, Milward K, Aliyu S, Makwana N. How clean are our stethoscopes and do we need to clean them? *J Infect* 2008;57:355–356.
14. Bernard L, Kereveur A, Durand D, et al. Bacterial contamination of hospital physicians' stethoscopes. *Infect Control Hosp Epidemiol* 1999;20:626–628.
15. Bukharie HA, Al-Zahrani H, Rubaish AM, Abdulmohsen MF. Bacterial contamination of stethoscopes. *J Fam Community Med* 2004;11:31–33.
16. Campos-Murguía A, León-Lara X, Muñoz JM, Macías AE, Álvarez JA. Stethoscopes as potential intrahospital carriers of pathogenic microorganisms. *Am J Infect Control* 2014;42:82–83.
17. Cohen HA, Amir J, Matalon A, Mayan R, Beni S, Barzilai A. Stethoscopes and otoscopes—a potential vector of infection? *Fam Pract* 1997;14:446–449.
18. Fafiora E, Bampalas VG, Lazarou N, et al. Bacterial contamination of medical devices in a Greek emergency department: impact of physicians' cleaning habits. *Am J Infect Control* 2014;42:807–809.
19. Fenelon L, Holcroft L, Waters N. Contamination of stethoscopes with MRSA and current disinfection practices. *J Hosp Infect* 2009;71:376–378.
20. Jones JS, Hoerle D, Riekse R. Stethoscopes: a potential vector of infection? *Ann Emerg Med* 1995;26:296–299.
21. Merlin MA, Wong ML, Pryor PW, et al. Prevalence of methicillin-resistant *Staphylococcus aureus* on the stethoscopes of emergency medical services providers. *Prehosp Emerg Care* 2009;13:71–74.
22. Núñez S, Moreno A, Green K, Villar J. The stethoscope in the emergency department: a vector of infection? *Epidemiol Infect* 2000;124:233–237.
23. Pandey A, Asthana AK, Tiwari R, Kumar L, Das A, Madan M. Physician accessories: doctor, what you carry is every patient's worry? *Indian J Pathol Microbiol* 2010;53:711–713.
24. Panhotra BR, Saxena AK, Al-Mulhim AS. Contaminated physician's stethoscope—a potential source of transmission of infection in the hospital. Need of frequent disinfection after use. *Saudi Med J* 2005;26:348–350.
25. Russell A, Secrest J, Schreeder C. Stethoscopes as a source of hospital-acquired methicillin-resistant *Staphylococcus aureus*. *J Perianesthesia Nurs* 2012;27:82–87.
26. Schroeder A, Schroeder MA, D'Amico F. What's growing on your stethoscope? (and what you can do about it). *J Fam Pract* 2009;58:404.
27. Sengupta S, Sirkar A, Shivananda PG. Stethoscopes and nosocomial infection. *Indian J Pediatr* 2000;67:197–199.
28. Smith MA, Mathewson JJ, Ulert IA, Scerpella EG, Ericsson CD. Contaminated stethoscopes revisited. *Arch Intern Med* 1996;156:82–84.
29. Shiferaw T, Beyene G, Kassa T, Sewunet T. Bacterial contamination, bacterial profile and antimicrobial susceptibility pattern of isolates from stethoscopes at Jimma University Specialized Hospital. *Ann Clin Microbiol Antimicrob* 2013;12:39.
30. Sood P, Mishra B, Mandal A. Potential infection hazards of stethoscopes. *J Indian Med Assoc* 2000;98:368–370.
31. Tang PH, Worster A, Srigley JA, Main CL. Examination of staphylococcal stethoscope contamination in the emergency department (pilot) study (EXSSCITED pilot study). *CJEM* 2011;13:239–244.
32. Uneke CJ, Ogbonna A, Oyibo PG, Onu CM. Bacterial contamination of stethoscopes used by health workers: public health implications. *J Infect Dev Ctries* 2010;4:436–441.
33. Whittington AM, Whitlow G, Hewson D, Thomas C, Brett SJ. Bacterial contamination of stethoscopes on the intensive care unit. *Anaesthesia* 2009;64:620–624.
34. Youngster I, Berkovitch M, Heyman E, Lazarovitch Z, Goldman M. The stethoscope as a vector of infectious diseases in the paediatric division. *Acta Paediatr* 2008;97:1253–1255.
35. Akinyemi KO, Atapu AD, Adetona OO, Coker AO. The potential role of mobile phones in the spread of bacterial infections. *J Infect Dev Ctries* 2009;3:628–632.
36. Beer D, Vandemere B, Brosnikoff C, Shokoples S, Rennie R, Forgie S. Bacterial contamination of health care workers' pagers and the efficacy of various disinfecting agents. *Pediatr Infect Dis J* 2006;25:1074–1075.
37. Borer A, Gilad J, Smolyakov R, et al. Cell phones and *Acinetobacter* transmission. *Emerg Infect Dis* 2005;11:1160–1161.
38. Braddy CM, Blair JE. Colonization of personal digital assistants used in a health care setting. *Am J Infect Control* 2005;33:230–232.
39. Brady RR, Fraser SF, Dunlop MG, Paterson-Brown S, Gibb AP. Bacterial contamination of mobile communication devices in the operative environment. *J Hosp Infect* 2007;66:397–398.
40. Brady RR, Wasson A, Stirling I, McAllister C, Damani NN. Is your phone bugged? The incidence of bacteria known to cause nosocomial infection on healthcare workers' mobile phones. *J Hosp Infect* 2006;62:123–125.
41. Datta P, Rani H, Chander J, Gupta V. Bacterial contamination of mobile phones of health care workers. *Indian J Med Microbiol* 2009;27:279–281.
42. Goldblatt JG, Krief I, Klonsky T, et al. Use of cellular telephones and transmission of pathogens by medical staff in New York and Israel. *Infect Control Hosp Epidemiol* 2007;28:500–503.
43. Hassoun A, Vellozzi EM, Smith MA. Colonization of personal digital assistants carried by healthcare professionals. *Infect Control Hosp Epidemiol* 2004;25:1000–1001.
44. Jayalakshmi J, Appalaraju B, Usha S. Cellphones as reservoirs of nosocomial pathogens. *J Assoc Physicians India* 2008;56:388–389.
45. Karabay O, Koçoglu E, Tahtaci M, et al. The role of mobile phones in the spread of bacteria associated with nosocomial infections. *J Infect Dev Ctries* 2007;1:72–73.
46. Kilic IH, Ozaslan M, Karagoz ID, Zer Y, Davutoglu V. The microbial colonisation of mobile phone used by healthcare staffs. *Pak J Biol Sci* 2009;12:882–884.
47. Lee YJ, Yoo CG, Lee CT, et al. Contamination rates between smart cell phones and non-smart cell phones of healthcare workers: bacterial contamination of smart phones. *J Hosp Med* 2013;8:144–147.
48. Namias N, Widrich J, Martinez OV, Cohn SM. Pathogenic bacteria on personal pagers. *Am J Infect Control* 2000;28:387–388.
49. Nwankwo EO, Ekwunife N, Mofolorunsho KC. Nosocomial pathogens associated with the mobile phones of healthcare workers in a hospital in Anyigba, Kogi state, Nigeria. *J Epidemiol Glob Health* 2014;4:135–140.
50. Ramesh J, Carter AO, Campbell MH, et al. Use of mobile phones by medical staff at Queen Elizabeth Hospital, Barbados: evidence for both benefit and harm. *J Hosp Infect* 2008;70:160–165.
51. Sadat-Ali M, Al-Omran AK, Azam Q, et al. Bacterial flora on cell phones of health care providers in a teaching institution. *Am J Infect Control* 2010;38:404–405.
52. Saxena S, Singh T, Agarwal H, Mehta G, Dutta R. Bacterial colonization of rings and cell phones carried by health-care

- providers: are these mobile bacterial zoos in the hospital? *Trop Doct* 2011;41:116–118.
53. Singh D, Kaur H, Gardner WG, Treen LB. Bacterial contamination of hospital papers. *Infect Control Hosp Epidemiol* 2002;23:274–276.
 54. Smith SJ, Knouse MC, Wasser T. Prevalence of bacterial pathogens on physician handheld computers. *J Clin Outcomes Manag* 2006;13:223–226.
 55. Srikanth P, Rajaram E, Sudharsanam S, et al. Mobile phones: emerging threat for infection control. *J Infect Prev* 2010;11:87–90.
 56. Tambekar DH, Gulhane PB, Dahikar SG, Dudhane MN. Nosocomial hazards of doctor's mobile phones in hospitals. *J Med Sci* 2008;8:73–76.
 57. Ulger F, Esen S, Dilek A, Yanik K, Gunaydin M, Leblebicioglu H. Are we aware how contaminated our mobile phones with nosocomial pathogens? *Ann Clin Microbiol Antimicrob* 2009;8:7.
 58. Ustun C, Cihangiroglu M. Health care workers' mobile phones: a potential cause of microbial cross-contamination between hospitals and community. *J Occup Environ Hyg* 2012;9:538–542.
 59. Walia SS, Manchanda A, Narang RS, Anup N, Singh B, Kahlon SS. Cellular telephone as reservoir of bacterial contamination: myth or fact. *J Clin Diagn Res* 2014;8:50–53.
 60. Loh W, Ng VV, Holton J. Bacterial flora on the white coats of medical students. *J Hosp Infect* 2000;45:65–68.
 61. Munoz-Price LS, Arheart KL, Mills JP, et al. Associations between bacterial contamination of health care workers' hands and contamination of white coats and scrubs. *Am J Infect Control* 2012;40:e245–e248.
 62. Treacle AM, Thom KM, Furuno JP, Strauss SM, Harris AD, Perencevich EN. Bacterial contamination of health care workers' white coats. *Am J Infect Control* 2009;37:101–105.
 63. Uneke CJ, Ijeoma PA. The potential for nosocomial infection transmission by white coats used by physicians in Nigeria: implications for improved patient-safety initiatives. *World Health Popul* 2010;11:44–54.
 64. Wiener-Well Y, Galuty M, Rudensky B, Schlesinger Y, Attias D, Yinnon DM. Nursing and physician attire as possible source of nosocomial infections. *Am J Infect Control* 2011;39:555–559.
 65. Ditchburn I. Should doctors wear ties? *J Hosp Infect* 2006;63:227–228.
 66. Koh KC, Husni S, Tan JE. High prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) on doctors' neckties. *Med J Malaysia* 2009;64:233–235.
 67. Lopez PJ, Ron O, Parthasarathy P, Soothill J, Spitz L. Bacterial counts from hospital doctors' ties are higher than those from shirts. *Am J Infect Control* 2009;37:79–80.
 68. McGovern B, Doyle E, Fenelon LE, FitzGerald SF. The necktie as a potential vector of infection: are doctors happy to do without? *J Hosp Infect* 2010;75:138–139.
 69. Bhat GK, Singhal L, Philip A, Jose T. Writing pens as fomites in hospital. *Indian J Med Microbiol* 2009;27:84–85.
 70. Datz C, Jungwirth A, Dusch H, Galvan G, Weiger T. What's on doctors' ball point pens? *Lancet* 1997;350:1824.
 71. Halton K, Arora V, Singh V, Ghantaji SS, Shah DN, Garey KW. Bacterial colonization on writing pens touched by healthcare professionals and hospitalized patients with and without cleaning the pen with alcohol-based hand sanitizing agent. *Clin Microbiol Infect* 2011;17:868–869.
 72. Wolfe DF, Sinnott S, Vossler JL, Przepiora J, Engbretson BG. Bacterial colonization of respiratory therapists' pens in the intensive care unit. *Respir Care* 2009;54:500–503.
 73. Feldman J, Feldman J, Feldman M. Women doctors' purses as an unrecognized fomite. *Del Med J* 2012;84:277–280.
 74. Gaspard P, Eschbach E, Gunther D, Gayet S, Bertrand X, Talon D. Methicillin-resistant *Staphylococcus aureus* contamination of healthcare workers' uniforms in long-term care facilities. *J Hosp Infect* 2009;71:170–175.
 75. Kotsanas D, Scott C, Gillespie EE, Korman TM, Stuart RL. What's hanging around your neck? Pathogenic bacteria on identity badges and lanyards. *Med J Aust* 2008;188:5.
 76. Ota K, Profitti R, Smaill F, Matlow AG, Smieja M. Identification badges: a potential fomite? *Can J Infect Control* 2007;22:162, 165–166.